

# Natural equivariant quantizations

Fabian Radoux

12th November 2008

## Introduction

- At the origin :  $qp \rightarrow QP$

$$qp \rightarrow \frac{1}{2}(QP + PQ)$$

with  $P = \partial_x$  ;  $Q = x$ .

## Introduction

- At the origin :  $qp \rightarrow QP$

$$qp \rightarrow \frac{1}{2}(QP + PQ)$$

with  $P = \partial_x$  ;  $Q = x$ .

Problems : It depends on the order

## Introduction

- At the origin :  $qp \rightarrow QP$

$$qp \rightarrow \frac{1}{2}(QP + PQ)$$

with  $P = \partial_x$  ;  $Q = x$ .

Problems : It depends on the order

It depends on the coordinates system

- Quantization :  $Q : \mathcal{S}(M) \mapsto \mathcal{D}_{\frac{1}{2}}(M)$

- Quantization :  $Q : \mathcal{S}(M) \mapsto \mathcal{D}_{\frac{1}{2}}(M)$

$Q$  : linear bijection

- Quantization :  $Q : \mathcal{S}(M) \mapsto \mathcal{D}_{\frac{1}{2}}(M)$

$Q$  : linear bijection

preserves the principal symbol

- Quantization :  $Q : \mathcal{S}(M) \mapsto \mathcal{D}_{\frac{1}{2}}(M)$

$Q$  : linear bijection

preserves the principal symbol

- $\#Q : Q(\Phi^* S) = \Phi^* Q(S) \forall$  local diffeomorphism  $\Phi$



- Quantization :  $Q : \mathcal{S}(M) \mapsto \mathcal{D}_{\frac{1}{2}}(M)$

$Q$  : linear bijection

preserves the principal symbol

- $\#Q : Q(\Phi^* S) = \Phi^* Q(S) \forall$  local diffeomorphism  $\Phi$   
 $\#Q : Q(L_X S) = L_X Q(S) \forall$  vector field  $X$

"Flat" case :

- $\exists Q : Q(\Phi_g^* S) = \Phi_g^* Q(S) \forall g \in G$

"Flat" case :

- $\exists Q : Q(\Phi_g^* S) = \Phi_g^* Q(S) \forall g \in G$   
 $\exists Q : Q(L_h^* S) = L_h^* Q(S) \forall h \in \mathfrak{g}$

"Flat" case :

- $\exists Q : Q(\Phi_g^* S) = \Phi_g^* Q(S) \forall g \in G$

$$\exists Q : Q(L_h^* S) = L_h^* Q(S) \forall h \in \mathfrak{g}$$

- $PGL(m+1, \mathbb{R})$  acts on  $\mathbb{R}P^m$

"Flat" case :

- $\exists Q : Q(\Phi_g^* S) = \Phi_g^* Q(S) \forall g \in G$

$$\exists Q : Q(L_h^* S) = L_h^* Q(S) \forall h \in \mathfrak{g}$$

- $PGL(m+1, \mathbb{R})$  acts on  $\mathbb{R}P^m$

$$X \in \mathfrak{sl}(m+1, \mathbb{R}) \mapsto X^* \text{ vector field on } \mathbb{R}^m$$

"Flat" case :

- $\exists Q : Q(\Phi_g^* S) = \Phi_g^* Q(S) \forall g \in G$

$$\exists Q : Q(L_h^* S) = L_h^* Q(S) \forall h \in \mathfrak{g}$$

- $PGL(m+1, \mathbb{R})$  acts on  $\mathbb{R}P^m$

$$X \in \mathfrak{sl}(m+1, \mathbb{R}) \mapsto X^* \text{ vector field on } \mathbb{R}^m$$

$$\exists Q : L_X Q(S) = Q(L_X S) \forall X \in \mathfrak{sl}(m+1, \mathbb{R})$$

"Flat" case :

- $\exists Q : Q(\Phi_g^* S) = \Phi_g^* Q(S) \forall g \in G$

$$\exists Q : Q(L_h^* S) = L_h^* Q(S) \forall h \in \mathfrak{g}$$

- $PGL(m+1, \mathbb{R})$  acts on  $\mathbb{R}P^m$

$$X \in \mathfrak{sl}(m+1, \mathbb{R}) \mapsto X^* \text{ vector field on } \mathbb{R}^m$$

$$\exists Q : L_X Q(S) = Q(L_X S) \forall X \in \mathfrak{sl}(m+1, \mathbb{R})$$

- $SO(p+1, q+1)$  acts on  $S^p \times S^q$

"Flat" case :

- $\exists Q : Q(\Phi_g^* S) = \Phi_g^* Q(S) \forall g \in G$

$$\exists Q : Q(L_h^* S) = L_h^* Q(S) \forall h \in \mathfrak{g}$$

- $PGL(m+1, \mathbb{R})$  acts on  $\mathbb{R}P^m$

$$X \in \mathfrak{sl}(m+1, \mathbb{R}) \mapsto X^* \text{ vector field on } \mathbb{R}^m$$

$$\exists Q : L_X Q(S) = Q(L_X S) \forall X \in \mathfrak{sl}(m+1, \mathbb{R})$$

- $SO(p+1, q+1)$  acts on  $S^p \times S^q$

$$X \in \mathfrak{so}(p+1, q+1) \mapsto X^* \text{ vector field on } \mathbb{R}^{p+q}$$



"Flat" case :

- $\exists Q : Q(\Phi_g^* S) = \Phi_g^* Q(S) \forall g \in G$

$$\exists Q : Q(L_h^* S) = L_h^* Q(S) \forall h \in \mathfrak{g}$$

- $PGL(m+1, \mathbb{R})$  acts on  $\mathbb{R}P^m$

$$X \in \mathfrak{sl}(m+1, \mathbb{R}) \mapsto X^* \text{ vector field on } \mathbb{R}^m$$

$$\exists Q : L_X Q(S) = Q(L_X S) \forall X \in \mathfrak{sl}(m+1, \mathbb{R})$$

- $SO(p+1, q+1)$  acts on  $S^p \times S^q$

$$X \in \mathfrak{so}(p+1, q+1) \mapsto X^* \text{ vector field on } \mathbb{R}^{p+q}$$

$$\exists Q : L_X Q(S) = Q(L_X S) \forall X \in \mathfrak{so}(p+1, q+1)$$

Introduction

Cartan fiber  
bundles and  
connections

The case of the  
densities

Other differential  
operators

Conformal case

Quantizations  
and foliations

- Method of the Casimir operator :

$$C : \mathcal{S}(\mathbb{R}^m) \mapsto \mathcal{S}(\mathbb{R}^m) \quad ; \quad \mathcal{C} : \mathcal{D}(\mathbb{R}^m) \mapsto \mathcal{D}(\mathbb{R}^m)$$

Introduction

Cartan fiber  
bundles and  
connections

The case of the  
densities

Other differential  
operators

Conformal case

Quantizations  
and foliations

- Method of the Casimir operator :

$$C : \mathcal{S}(\mathbb{R}^m) \mapsto \mathcal{S}(\mathbb{R}^m) \quad ; \quad \mathcal{C} : \mathcal{D}(\mathbb{R}^m) \mapsto \mathcal{D}(\mathbb{R}^m)$$

Condition :  $\exists Q$  such that if  $C(S) = \alpha S$ , then

Introduction

Cartan fiber  
bundles and  
connections

The case of the  
densities

Other differential  
operators

Conformal case

Quantizations  
and foliations

- Method of the Casimir operator :

$$\mathcal{C} : \mathcal{S}(\mathbb{R}^m) \mapsto \mathcal{S}(\mathbb{R}^m) \quad ; \quad \mathcal{C} : \mathcal{D}(\mathbb{R}^m) \mapsto \mathcal{D}(\mathbb{R}^m)$$

Condition :  $\exists Q$  such that if  $\mathcal{C}(S) = \alpha S$ , then

$$\mathcal{C}(Q(S)) = \alpha Q(S)$$

- Method of the Casimir operator :

$$\mathcal{C} : \mathcal{S}(\mathbb{R}^m) \mapsto \mathcal{S}(\mathbb{R}^m) \quad ; \quad \mathcal{C} : \mathcal{D}(\mathbb{R}^m) \mapsto \mathcal{D}(\mathbb{R}^m)$$

Condition :  $\exists Q$  such that if  $\mathcal{C}(S) = \alpha S$ , then

$$\mathcal{C}(Q(S)) = \alpha Q(S) \text{ and } \sigma(Q(S)) = S$$

"Curved" case :

- $Q(\nabla) : \mathcal{S}^3(M) \mapsto \mathcal{D}^3(M)$

$$Q(\nabla) = Q(\nabla') \text{ if } \nabla' = \nabla + \alpha \vee id$$

"Curved" case :

- $Q(\nabla) : \mathcal{S}^3(M) \mapsto \mathcal{D}^3(M)$

$$Q(\nabla) = Q(\nabla') \text{ if } \nabla' = \nabla + \alpha \vee id$$

- $Q(g) : \mathcal{S}^2(M) \mapsto \mathcal{D}^2(M)$

$$Q(g) = Q(g') \text{ if } g = fg' \text{ with } f > 0$$

Introduction

Cartan fiber  
bundles and  
connections

The case of the  
densities

Other differential  
operators

Conformal case

Quantizations  
and foliations

- Conjecture :  $Q(g) : \mathcal{S}(M) \mapsto \mathcal{D}(M)$



## Introduction

Cartan fiber  
bundles and  
connections

The case of the  
densities

Other differential  
operators

Conformal case

Quantizations  
and foliations

- Conjecture :  $Q(g) : \mathcal{S}(M) \mapsto \mathcal{D}(M)$  natural

- Conjecture :  $Q(g) : \mathcal{S}(M) \mapsto \mathcal{D}(M)$  natural and
$$Q(g) = Q(g') \text{ if } g = fg' \text{ with } f > 0$$

- Conjecture :  $Q(g) : \mathcal{S}(M) \mapsto \mathcal{D}(M)$  natural and

$$Q(g) = Q(g') \text{ if } g = fg' \text{ with } f > 0$$

$$\varphi_t^* Q(g^0)(S) = Q(\varphi_t^* g^0)(\varphi_t^* S)$$

- Conjecture :  $Q(g) : \mathcal{S}(M) \mapsto \mathcal{D}(M)$  natural and

$$Q(g) = Q(g') \text{ if } g = fg' \text{ with } f > 0$$

$$\varphi_t^* Q(g^0)(S) = Q(\varphi_t^* g^0)(\varphi_t^* S)$$

$$L_X Q(g^0)(S) = Q(g^0)(L_X S) \quad \forall X \in \mathfrak{so}(p+1, q+1)$$

- Conjecture :  $Q(g) : \mathcal{S}(M) \mapsto \mathcal{D}(M)$  natural and

$$Q(g) = Q(g') \text{ if } g = fg' \text{ with } f > 0$$

$$\varphi_t^* Q(g^0)(S) = Q(\varphi_t^* g^0)(\varphi_t^* S)$$

$$L_X Q(g^0)(S) = Q(g^0)(L_X S) \quad \forall X \in \mathfrak{so}(p+1, q+1)$$

- Conjecture :  $Q(\nabla) : \mathcal{S}(M) \mapsto \mathcal{D}(M)$

- Conjecture :  $Q(g) : \mathcal{S}(M) \mapsto \mathcal{D}(M)$  natural and

$$Q(g) = Q(g') \text{ if } g = fg' \text{ with } f > 0$$

$$\varphi_t^* Q(g^0)(S) = Q(\varphi_t^* g^0)(\varphi_t^* S)$$

$$L_X Q(g^0)(S) = Q(g^0)(L_X S) \quad \forall X \in \mathfrak{so}(p+1, q+1)$$

- Conjecture :  $Q(\nabla) : \mathcal{S}(M) \mapsto \mathcal{D}(M)$  natural

- Conjecture :  $Q(g) : \mathcal{S}(M) \mapsto \mathcal{D}(M)$  natural and

$$Q(g) = Q(g') \text{ if } g = fg' \text{ with } f > 0$$

$$\varphi_t^* Q(g^0)(S) = Q(\varphi_t^* g^0)(\varphi_t^* S)$$

$$L_X Q(g^0)(S) = Q(g^0)(L_X S) \quad \forall X \in \mathfrak{so}(p+1, q+1)$$

- Conjecture :  $Q(\nabla) : \mathcal{S}(M) \mapsto \mathcal{D}(M)$  natural and

$$Q(\nabla) = Q(\nabla') \text{ if } \nabla' = \nabla + \alpha \vee id$$

- Conjecture :  $Q(g) : \mathcal{S}(M) \mapsto \mathcal{D}(M)$  natural and

$$Q(g) = Q(g') \text{ if } g = fg' \text{ with } f > 0$$

$$\varphi_t^* Q(g^0)(S) = Q(\varphi_t^* g^0)(\varphi_t^* S)$$

$$L_X Q(g^0)(S) = Q(g^0)(L_X S) \quad \forall X \in \mathfrak{so}(p+1, q+1)$$

- Conjecture :  $Q(\nabla) : \mathcal{S}(M) \mapsto \mathcal{D}(M)$  natural and

$$Q(\nabla) = Q(\nabla') \text{ if } \nabla' = \nabla + \alpha \vee id$$

$$\phi_t^* Q(\nabla^0)(S) = Q(\phi_t^* \nabla^0)(\phi_t^* S)$$



- Conjecture :  $Q(g) : \mathcal{S}(M) \mapsto \mathcal{D}(M)$  natural and

$$Q(g) = Q(g') \text{ if } g = fg' \text{ with } f > 0$$

$$\varphi_t^* Q(g^0)(S) = Q(\varphi_t^* g^0)(\varphi_t^* S)$$

$$L_X Q(g^0)(S) = Q(g^0)(L_X S) \quad \forall X \in \mathfrak{so}(p+1, q+1)$$

- Conjecture :  $Q(\nabla) : \mathcal{S}(M) \mapsto \mathcal{D}(M)$  natural and

$$Q(\nabla) = Q(\nabla') \text{ if } \nabla' = \nabla + \alpha \vee id$$

$$\phi_t^* Q(\nabla^0)(S) = Q(\phi_t^* \nabla^0)(\phi_t^* S)$$

$$L_X Q(\nabla^0)(S) = Q(\nabla^0)(L_X S) \quad \forall X \in \mathfrak{sl}(m+1, \mathbb{R})$$

# Natural equivariant quantizations

Fabian Radoux

## Introduction

Cartan fiber  
bundles and  
connections

The case of the  
densities

Other differential  
operators

Conformal case

Quantizations  
and foliations

- M. Bordemann method :

Natural  
equivariant  
quantizations

Fabian Radoux

Introduction

Cartan fiber  
bundles and  
connections

The case of the  
densities

Other differential  
operators

Conformal case

Quantizations  
and foliations

- M. Bordemann method :  $M \mapsto \tilde{M}$ ;

- M. Bordemann method :  $M \mapsto \tilde{M}$ ;

$$\nabla \mapsto \tilde{\nabla};$$

- M. Bordemann method :  $M \mapsto \tilde{M}$ ;

$$\nabla \mapsto \tilde{\nabla};$$

$$S \mapsto \tilde{S};$$

- M. Bordemann method :  $M \mapsto \tilde{M}$ ;

$$\nabla \mapsto \tilde{\nabla};$$

$$S \mapsto \tilde{S};$$

$$Q(\widetilde{\nabla})(\tilde{S})(f) = \tau(\tilde{\nabla})(\tilde{S})(\tilde{f})$$

# Natural equivariant quantizations

Fabian Radoux

## Introduction

Cartan fiber  
bundles and  
connections

The case of the  
densities

Other differential  
operators

Conformal case

Quantizations  
and foliations

- Questions :

Natural  
equivariant  
quantizations

Fabian Radoux

Introduction

Cartan fiber  
bundles and  
connections

The case of the  
densities

Other differential  
operators

Conformal case

Quantizations  
and foliations

- Questions : Critical values of  $\delta$



- Questions : Critical values of  $\delta$   
Explicit formula

- Questions : Critical values of  $\delta$

Explicit formula

Uniqueness

- Questions : Critical values of  $\delta$

Explicit formula

Uniqueness

Other differential operators

Introduction

Cartan fiber  
bundles and  
connections

The case of the  
densities

Other differential  
operators

Conformal case

Quantizations  
and foliations

- Questions : Critical values of  $\delta$

Explicit formula

Uniqueness

Other differential operators

Conformal case

- Cartan fiber bundles and connections

- Cartan fiber bundles and connections

$$[\nabla] \rightarrow P \mapsto M$$

- Cartan fiber bundles and connections

$$[\nabla] \rightarrow P \mapsto M$$

$$[\nabla] \rightarrow \omega : TP \mapsto \mathfrak{g}$$

- Cartan fiber bundles and connections

$$[\nabla] \rightarrow P \mapsto M$$

$$[\nabla] \rightarrow \omega : TP \mapsto \mathfrak{g}$$

$$\omega_u : T_u P \mapsto \mathfrak{g} \text{ bijection } \forall u \in P$$



- Cartan fiber bundles and connections

$$[\nabla] \rightarrow P \mapsto M$$

$$[\nabla] \rightarrow \omega : TP \mapsto \mathfrak{g}$$

$$\omega_u : T_u P \mapsto \mathfrak{g} \text{ bijection } \forall u \in P$$

$$T \rightarrow p^* T \in \mathcal{C}^\infty(P, V)_H$$

- Cartan fiber bundles and connections

$$[\nabla] \rightarrow P \mapsto M$$

$$[\nabla] \rightarrow \omega : TP \mapsto \mathfrak{g}$$

$$\omega_u : T_u P \mapsto \mathfrak{g} \text{ bijection } \forall u \in P$$

$$T \rightarrow p^* T \in \mathcal{C}^\infty(P, V)_H$$

$$\text{with } H = G_0 \rtimes G_1, \mathfrak{h} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$$

- Cartan fiber bundles and connections

$$[\nabla] \rightarrow P \mapsto M$$

$$[\nabla] \rightarrow \omega : TP \mapsto \mathfrak{g}$$

$$\omega_u : T_u P \mapsto \mathfrak{g} \text{ bijection } \forall u \in P$$

$$T \rightarrow p^* T \in \mathcal{C}^\infty(P, V)_H$$

$$\text{with } H = G_0 \rtimes G_1, \mathfrak{h} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$$

$$\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{h}, \mathfrak{g}_{-1} \cong \mathbb{R}^m$$

- Cartan fiber bundles and connections

$$[\nabla] \rightarrow P \mapsto M$$

$$[\nabla] \rightarrow \omega : TP \mapsto \mathfrak{g}$$

$$\omega_u : T_u P \mapsto \mathfrak{g} \text{ bijection } \forall u \in P$$

$$T \rightarrow p^* T \in \mathcal{C}^\infty(P, V)_H$$

$$\text{with } H = G_0 \rtimes G_1, \mathfrak{h} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$$

$$\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{h}, \mathfrak{g}_{-1} \cong \mathbb{R}^m$$

$$\omega \rightarrow \nabla^\omega(e_i) = L_{\omega^{-1}(e_i)}, e_i \in \mathfrak{g}_{-1}$$

## • Cartan fiber bundles and connections

$$[\nabla] \rightarrow P \mapsto M$$

$$[\nabla] \rightarrow \omega : TP \mapsto \mathfrak{g}$$

$$\omega_u : T_u P \mapsto \mathfrak{g} \text{ bijection } \forall u \in P$$

$$T \rightarrow p^* T \in \mathcal{C}^\infty(P, V)_H$$

$$\text{with } H = G_0 \rtimes G_1, \mathfrak{h} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$$

$$\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{h}, \mathfrak{g}_{-1} \cong \mathbb{R}^m$$

$$\omega \rightarrow \nabla^\omega(e_i) = L_{\omega^{-1}(e_i)}, e_i \in \mathfrak{g}_{-1}$$

$$f \text{ } G_0\text{-equivariant} \Rightarrow \nabla^\omega f$$

$G_0$ -equivariant

## • Cartan fiber bundles and connections

$$[\nabla] \rightarrow P \mapsto M$$

$$[\nabla] \rightarrow \omega : TP \mapsto \mathfrak{g}$$

$$\omega_u : T_u P \mapsto \mathfrak{g} \text{ bijection } \forall u \in P$$

$$T \rightarrow p^* T \in \mathcal{C}^\infty(P, V)_H$$

$$\text{with } H = G_0 \rtimes G_1, \mathfrak{h} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$$

$$\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{h}, \mathfrak{g}_{-1} \cong \mathbb{R}^m$$

$$\omega \rightarrow \nabla^\omega(e_i) = L_{\omega^{-1}(e_i)}, e_i \in \mathfrak{g}_{-1}$$

$$f \text{ } G_0\text{-equivariant} \Rightarrow \nabla^\omega f$$

$G_0$ -equivariant

$$f \text{ } G_1\text{-equivariant} \not\Rightarrow \nabla^\omega f$$

$G_1$ -equivariant

- The case of the densities (P. Mathonet, R.) :

- The case of the densities (P. Mathonet, R.) :

$$S \rightarrow p^* S \in \mathcal{C}^\infty(P, S_\delta^k(\mathbb{R}^m))_H$$



- The case of the densities (P. Mathonet, R.) :

$$S \rightarrow p^* S \in \mathcal{C}^\infty(P, S_\delta^k(\mathbb{R}^m))_H$$

$$f \rightarrow p^* f \in \mathcal{C}^\infty(P, \Delta^\lambda(\mathbb{R}^m))_H$$

- The case of the densities (P. Mathonet, R.) :

$$S \rightarrow p^* S \in \mathcal{C}^\infty(P, S_\delta^k(\mathbb{R}^m))_H$$

$$f \rightarrow p^* f \in \mathcal{C}^\infty(P, \Delta^\lambda(\mathbb{R}^m))_H$$

$$\omega \rightarrow \text{Div}^\omega = \sum_i i(\epsilon^i) L_{\omega^{-1}(e_i)}$$

- The case of the densities (P. Mathonet, R.) :

$$S \rightarrow p^* S \in \mathcal{C}^\infty(P, S_\delta^k(\mathbb{R}^m))_H$$

$$f \rightarrow p^* f \in \mathcal{C}^\infty(P, \Delta^\lambda(\mathbb{R}^m))_H$$

$$\omega \rightarrow \text{Div}^\omega = \sum_i i(\epsilon^i) L_{\omega^{-1}(\epsilon_i)}$$

$$\text{Condition : } L_{h^*} Q(p^* S)(p^* f) = 0 \quad \forall h \in \mathfrak{g}_1$$

- The case of the densities (P. Mathonet, R.) :

$$S \rightarrow p^* S \in \mathcal{C}^\infty(P, S_\delta^k(\mathbb{R}^m))_H$$

$$f \rightarrow p^* f \in \mathcal{C}^\infty(P, \Delta^\lambda(\mathbb{R}^m))_H$$

$$\omega \rightarrow \text{Div}^\omega = \sum_i i(\epsilon^i) L_{\omega^{-1}(e_i)}$$

Condition :  $L_{h^*} Q(p^* S)(p^* f) = 0 \ \forall h \in \mathfrak{g}_1$

$\langle p^* S, \nabla_s^{\omega^k} p^* f \rangle$  not  $G_1$ -equivariant !

- The case of the densities (P. Mathonet, R.) :

$$S \rightarrow p^* S \in \mathcal{C}^\infty(P, S_\delta^k(\mathbb{R}^m))_H$$

$$f \rightarrow p^* f \in \mathcal{C}^\infty(P, \Delta^\lambda(\mathbb{R}^m))_H$$

$$\omega \rightarrow \text{Div}^\omega = \sum_i i(\epsilon^i) L_{\omega^{-1}(\epsilon_i)}$$

Condition :  $L_{h^*} Q(p^* S)(p^* f) = 0 \ \forall h \in \mathfrak{g}_1$

$\langle p^* S, \nabla_s^{\omega^k} p^* f \rangle$  not  $G_1$ -equivariant !

One adds terms of lower orders in  $p^* f \dots$

- The case of the densities (P. Mathonet, R.) :

$$S \rightarrow p^* S \in \mathcal{C}^\infty(P, S_\delta^k(\mathbb{R}^m))_H$$

$$f \rightarrow p^* f \in \mathcal{C}^\infty(P, \Delta^\lambda(\mathbb{R}^m))_H$$

$$\omega \rightarrow \text{Div}^\omega = \sum_i i(\epsilon^i) L_{\omega^{-1}(\epsilon_i)}$$

Condition :  $L_{h^*} Q(p^* S)(p^* f) = 0 \ \forall h \in \mathfrak{g}_1$

$\langle p^* S, \nabla_s^{\omega^k} p^* f \rangle$  not  $G_1$ -equivariant !

One adds terms of lower orders in  $p^* f \dots$

One finds then :

$$Q_M(\nabla, S)(f) = p^{*-1} \left( \sum_{l=0}^k C_{k,l} \langle \text{Div}^{\omega^l} p^* S, \nabla_s^{\omega^{k-l}} p^* f \rangle \right),$$

$$\text{with } C_{k,l} = \frac{(\lambda + \frac{k-1}{m+1}) \dots (\lambda + \frac{k-l}{m+1})}{\gamma_{2k-1} \dots \gamma_{2k-l}} \binom{k}{l}, \quad \forall l \geq 1, \quad C_{k,0} = 1$$

- Other differential operators (P. Mathonet, R.) : method of the Casimir operator :

- Other differential operators (P. Mathonet, R.) : method of the Casimir operator :

"Flat" case

"Curved" case



- Other differential operators (P. Mathonet, R.) : method of the Casimir operator :

"Flat" case

Affine quantization  $Q_{Aff} :$   
 $\partial_i$

"Curved" case

"Affine" quantization  $Q_\omega :$   
 $L_{\omega^{-1}(e_i)}$

- Other differential operators (P. Mathonet, R.) : method of the Casimir operator :

"Flat" case

Affine quantization  $Q_{Aff} :$   
 $\partial_i$

Application  $\gamma :$

$$\mathcal{L}_{X^h} Q_{Aff}(S)(f) = \\ Q_{Aff}((L_{X^h} + \gamma(h))S)(f)$$

"Curved" case

"Affine" quantization  $Q_\omega :$   
 $L_{\omega^{-1}(e_i)}$

Application  $\gamma :$

$$\mathcal{L}_{h^*} Q_\omega(S)(f) = \\ Q_\omega((L_{h^*} + \gamma(h))S)(f), \quad h \in \mathfrak{g}_1$$

• Other differential operators (P. Mathonet, R.) : method of the Casimir operator :

"Flat" case

Affine quantization  $Q_{Aff} :$   
 $\partial_i$

Application  $\gamma :$

$$\mathcal{L}_{X^h} Q_{Aff}(S)(f) =$$

$$Q_{Aff}((L_{X^h} + \gamma(h))S)(f)$$

Casimirs  $C$  and  $\mathcal{C}$

$$\partial_i$$

"Curved" case

"Affine" quantization  $Q_\omega :$   
 $L_{\omega^{-1}(e_i)}$

Application  $\gamma :$

$$\mathcal{L}_{h^*} Q_\omega(S)(f) =$$

$$Q_\omega((L_{h^*} + \gamma(h))S)(f), \quad h \in \mathfrak{g}_1$$

"Casimirs"  $C^\omega$  and  $\mathcal{C}^\omega$

$$L_{\omega^{-1}(e_i)}$$

Quantization :

$Q_{Aff}(Q(S))$ ,  $Q(S)$  such that

if  $C(S) = \alpha S$ , then

$\mathcal{C}(Q(S)) = \alpha Q(S)$  and

"head" of  $Q(S) = S$

Quantization :

$Q_{\omega}(Q(S))$ ,  $Q(S)$  such that

if  $C^{\omega}(S) = \alpha S$ , then

$\mathcal{C}^{\omega}(Q(S)) = \alpha Q(S)$  and

"head" of  $Q(S) = S$

Quantization :

$Q_{Aff}(Q(S))$ ,  $Q(S)$  such that

if  $C(S) = \alpha S$ , then

$\mathcal{C}(Q(S)) = \alpha Q(S)$  and

"head" of  $Q(S) = S$

Then :

$$\mathcal{L} \circ Q = Q \circ L$$

because

$$[\mathcal{C}, \mathcal{L}] = 0 \text{ and } [C, L] = 0$$

Quantization :

$Q_{\omega}(Q(S))$ ,  $Q(S)$  such that

if  $C^{\omega}(S) = \alpha S$ , then

$\mathcal{C}^{\omega}(Q(S)) = \alpha Q(S)$  and

"head" of  $Q(S) = S$

Then :

$$(L_{h^*} + \gamma(h)) \circ Q = Q \circ L_{h^*}$$

because  $[\mathcal{C}^{\omega}, L_{h^*} + \gamma(h)] = 0$

$$\text{and } [C^{\omega}, L_{h^*}] = 0$$

Quantization :

$Q_{Aff}(Q(S))$ ,  $Q(S)$  such that

if  $C(S) = \alpha S$ , then

$\mathcal{C}(Q(S)) = \alpha Q(S)$  and

"head" of  $Q(S) = S$

Then :

$$\mathcal{L} \circ Q = Q \circ L$$

because

$$[\mathcal{C}, \mathcal{L}] = 0 \text{ and } [C, L] = 0$$

• Conclusion : "Flat" case  
 $\partial_i$

Quantization :

$Q_\omega(Q(S))$ ,  $Q(S)$  such that

if  $C^\omega(S) = \alpha S$ , then

$\mathcal{C}^\omega(Q(S)) = \alpha Q(S)$  and

"head" of  $Q(S) = S$

Then :

$$(L_{h^*} + \gamma(h)) \circ Q = Q \circ L_{h^*}$$

because  $[\mathcal{C}^\omega, L_{h^*} + \gamma(h)] = 0$

and  $[\mathcal{C}^\omega, L_{h^*}] = 0$

"Curved" case  
 $L_{\omega^{-1}(e_i)}$

Introduction

Cartan fiber  
bundles and  
connections

The case of the  
densities

Other differential  
operators

**Conformal case**

Quantizations  
and foliations

- Conformal case (P. Mathonet, R.) :

- Conformal case (P. Mathonet, R.) :

"Affine" quantization  $Q'_\omega : Q_\omega$  "not symmetrized"



- Conformal case (P. Mathonet, R.) :

"Affine" quantization  $Q'_\omega : Q_\omega$  "not symmetrized"

Application  $\gamma' : \gamma$  "not symmetric"

- Conformal case (P. Mathonet, R.) :

"Affine" quantization  $Q'_\omega : Q_\omega$  "not symmetrized"

Application  $\gamma' : \gamma$  "not symmetric"

$$\mathcal{L}_{h^*} Q'_\omega(S)(f) = Q'_\omega((L_{h^*} + \gamma'(h))S)(f), \quad h \in \mathfrak{g}_1$$

- Conformal case (P. Mathonet, R.) :

"Affine" quantization  $Q'_\omega : Q_\omega$  "not symmetrized"

Application  $\gamma' : \gamma$  "not symmetric"

$$\mathcal{L}_{h^*} Q'_\omega(S)(f) = Q'_\omega((L_{h^*} + \gamma'(h))S)(f), \quad h \in \mathfrak{g}_1$$

"Casimirs"  $\mathcal{C}^{\omega'} = \mathcal{C}^\omega$  and  $\mathcal{C}^{\omega'}$

• Conformal case (P. Mathonet, R.) :

"Affine" quantization  $Q'_\omega : Q_\omega$  "not symmetrized"

Application  $\gamma' : \gamma$  "not symmetric"

$$\mathcal{L}_{h^*} Q'_\omega(S)(f) = Q'_\omega((L_{h^*} + \gamma'(h))S)(f), \quad h \in \mathfrak{g}_1$$

"Casimirs"  $\mathcal{C}^{\omega'}$  " = "  $\mathcal{C}^\omega$  and  $\mathcal{C}^{\omega'}$

Quantization :  $Q'_\omega(Q'(S))$ ,  $Q'(S)$  such that if  $\mathcal{C}^{\omega'}(S) = \alpha S$ ,  
then  $\mathcal{C}^{\omega'}(Q'(S)) = \alpha Q'(S)$  and "head" of  $Q'(S) = S$

• Conformal case (P. Mathonet, R.) :

"Affine" quantization  $Q'_\omega : Q_\omega$  "not symmetrized"

Application  $\gamma' : \gamma$  "not symmetric"

$$\mathcal{L}_{h^*} Q'_\omega(S)(f) = Q'_\omega((L_{h^*} + \gamma'(h))S)(f), \quad h \in \mathfrak{g}_1$$

"Casimirs"  $\mathcal{C}^{\omega'}$  " = "  $\mathcal{C}^\omega$  and  $\mathcal{C}^{\omega'}$

Quantization :  $Q'_\omega(Q'(S))$ ,  $Q'(S)$  such that if  $\mathcal{C}^{\omega'}(S) = \alpha S$ , then  $\mathcal{C}^{\omega'}(Q'(S)) = \alpha Q'(S)$  and "head" of  $Q'(S) = S$

Then :  $(L_{h^*} + \gamma'(h)) \circ Q' = Q' \circ L_{h^*}$  because  
 $[\mathcal{C}^{\omega'}, L_{h^*} + \gamma'(h)] = 0$  and  $[\mathcal{C}^{\omega'}, L_{h^*}] = 0$

Natural  
equivariant  
quantizations

Fabian Radoux

Introduction

Cartan fiber  
bundles and  
connections

The case of the  
densities

Other differential  
operators

Conformal case

Quantizations  
and foliations

Remark : allows to find natural applications

$Q : \{\text{reductions of } P^2M \text{ to } H\} \rightarrow \{\text{quantizations on } M\},$

where  $P^2M$  is the second order frame bundle and where  $H$  is a group linked to an AHS structure

- Quantizations and foliations (N. Poncin, R. Wolak, R.) :
- Quantize  $M/G \leftrightarrow$  Quantize  $M/\bar{\mathcal{F}}$

- Quantizations and foliations (N. Poncin, R. Wolak, R.) :
- Quantize  $M/G \leftrightarrow$  Quantize  $M/\bar{\mathcal{F}}$

$$\begin{array}{ccc} Q_M(\nabla_M)(S_M) : M & \longrightarrow & M \\ & \pi \downarrow & \downarrow \pi \\ Q_{M/\bar{\mathcal{F}}}(\nabla_{M/\bar{\mathcal{F}}})(S_{M/\bar{\mathcal{F}}}) : M/\bar{\mathcal{F}} & \longrightarrow & M/\bar{\mathcal{F}} \end{array}$$



- Quantizations and foliations (N. Poncin, R. Wolak, R.) :
- Quantize  $M/G \leftrightarrow$  Quantize  $M/\bar{\mathcal{F}}$

$$\begin{array}{ccc}
 Q_M(\nabla_M)(S_M) : M & \longrightarrow & M \\
 & \pi \downarrow & \downarrow \pi \\
 Q_{M/\bar{\mathcal{F}}}(\nabla_{M/\bar{\mathcal{F}}})(S_{M/\bar{\mathcal{F}}}) : M/\bar{\mathcal{F}} & \longrightarrow & M/\bar{\mathcal{F}}
 \end{array}$$

Quantization commutes with reduction :

$$\pi(Q_M(\nabla_M)(S_M)) = Q_{M/\bar{\mathcal{F}}}(\pi\nabla_M)(\pi S_M)$$

- Quantizations and foliations (N. Poncin, R. Wolak, R.) :
- Quantize  $M/G \leftrightarrow$  Quantize  $M/\bar{\mathcal{F}}$

$$\begin{array}{ccc}
 Q_M(\nabla_M)(S_M) : M & \longrightarrow & M \\
 & \pi \downarrow & \downarrow \pi \\
 Q_{M/\bar{\mathcal{F}}}(\nabla_{M/\bar{\mathcal{F}}})(S_{M/\bar{\mathcal{F}}}) : M/\bar{\mathcal{F}} & \longrightarrow & M/\bar{\mathcal{F}}
 \end{array}$$

Quantization commutes with reduction :

$$\pi(Q_M(\nabla_M)(S_M)) = Q_{M/\bar{\mathcal{F}}}(\pi\nabla_M)(\pi S_M)$$

$\nabla_M, S_M, Q_M(\nabla_M)(S_M) \leftrightarrow$  adapted objects

- Quantizations and foliations (N. Poncin, R. Wolak, R.) :

- Quantize  $M/G \leftrightarrow$  Quantize  $M/\bar{\mathcal{F}}$

$$\begin{array}{ccc}
 Q_M(\nabla_M)(S_M) : M & \longrightarrow & M \\
 & \pi \downarrow & \downarrow \pi \\
 Q_{M/\bar{\mathcal{F}}}(\nabla_{M/\bar{\mathcal{F}}})(S_{M/\bar{\mathcal{F}}}) : M/\bar{\mathcal{F}} & \longrightarrow & M/\bar{\mathcal{F}}
 \end{array}$$

Quantization commutes with reduction :

$$\pi(Q_M(\nabla_M)(S_M)) = Q_{M/\bar{\mathcal{F}}}(\pi\nabla_M)(\pi S_M)$$

$\nabla_M, S_M, Q_M(\nabla_M)(S_M) \leftrightarrow$  adapted objects

$\nabla_{M/\bar{\mathcal{F}}}, S_{M/\bar{\mathcal{F}}}, Q_{M/\bar{\mathcal{F}}}(\nabla_{M/\bar{\mathcal{F}}})(S_{M/\bar{\mathcal{F}}}) \approx$  foliated objects

Adapted vector field :  $X$  such that  $[X, Y] \in T\mathcal{F} \forall Y \in T\mathcal{F}$   
( $\rightarrow Vect_{\mathcal{F}}(M)$ )

Adapted vector field :  $X$  such that  $[X, Y] \in T\mathcal{F} \forall Y \in T\mathcal{F}$   
( $\rightarrow Vect_{\mathcal{F}}(M)$ )

In adapted coordinates  $(x, y)$  :

$$X = \sum_i X^i(x, y) \partial_{x^i} + \sum_i X^i(y) \partial_{y^i}$$

Adapted vector field :  $X$  such that  $[X, Y] \in T\mathcal{F} \forall Y \in T\mathcal{F}$   
( $\rightarrow Vect_{\mathcal{F}}(M)$ )

In adapted coordinates  $(x, y)$  :

$$X = \sum_i X^i(x, y) \partial_{x^i} + \sum_i X^i(y) \partial_{y^i}$$

Foliated vector field :  $[X]$  with  $X$  adapted and with  
 $[ \ ] : TM \rightarrow TM/T\mathcal{F}$  ( $\rightarrow Vect(M, \mathcal{F})$ )

Adapted vector field :  $X$  such that  $[X, Y] \in T\mathcal{F} \forall Y \in T\mathcal{F}$   
( $\rightarrow Vect_{\mathcal{F}}(M)$ )

In adapted coordinates  $(x, y)$  :

$$X = \sum_i X^i(x, y) \partial_{x^i} + \sum_i X^i(y) \partial_{y^i}$$

Foliated vector field :  $[X]$  with  $X$  adapted and with  
 $[ \ ] : TM \rightarrow TM/T\mathcal{F}$  ( $\rightarrow Vect(M, \mathcal{F})$ )

In adapted coordinates  $(x, y)$  :  $[X] = \sum_i X^i(y) [\partial_{y^i}]$

Adapted vector field :  $X$  such that  $[X, Y] \in T\mathcal{F} \forall Y \in T\mathcal{F}$   
( $\rightarrow Vect_{\mathcal{F}}(M)$ )

In adapted coordinates  $(x, y)$  :

$$X = \sum_i X^i(x, y) \partial_{x^i} + \sum_i X^i(y) \partial_{y^i}$$

Foliated vector field :  $[X]$  with  $X$  adapted and with  
 $[ \ ] : TM \rightarrow TM/T\mathcal{F}$  ( $\rightarrow Vect(M, \mathcal{F})$ )

In adapted coordinates  $(x, y)$  :  $[X] = \sum_i X^i(y) [\partial_{y^i}]$

Projection :  $\sum_i X^i(x, y) \partial_{x^i} + \sum_i X^i(y) \partial_{y^i} \mapsto \sum_i X^i(y) [\partial_{y^i}]$



Adapted connection :

$$\begin{aligned} \nabla_{\mathcal{F}} : Vect_{\mathcal{F}}(M) \times Vect_{\mathcal{F}}(M) &\rightarrow Vect_{\mathcal{F}}(M) \\ &: Vect_{\mathcal{F}}(M) \times T\mathcal{F} \rightarrow T\mathcal{F} \end{aligned}$$

Adapted connection :

$$\begin{aligned}\nabla_{\mathcal{F}} : \text{Vect}_{\mathcal{F}}(M) \times \text{Vect}_{\mathcal{F}}(M) &\rightarrow \text{Vect}_{\mathcal{F}}(M) \\ &: \text{Vect}_{\mathcal{F}}(M) \times T\mathcal{F} \rightarrow T\mathcal{F}\end{aligned}$$

Foliated connection :

$$\nabla(\mathcal{F}) : \text{Vect}(M, \mathcal{F}) \times \text{Vect}(M, \mathcal{F}) \rightarrow \text{Vect}(M, \mathcal{F})$$

Adapted connection :

$$\begin{aligned}\nabla_{\mathcal{F}} : \text{Vect}_{\mathcal{F}}(M) \times \text{Vect}_{\mathcal{F}}(M) &\rightarrow \text{Vect}_{\mathcal{F}}(M) \\ &: \text{Vect}_{\mathcal{F}}(M) \times T\mathcal{F} \rightarrow T\mathcal{F}\end{aligned}$$

Foliated connection :

$$\nabla(\mathcal{F}) : \text{Vect}(M, \mathcal{F}) \times \text{Vect}(M, \mathcal{F}) \rightarrow \text{Vect}(M, \mathcal{F})$$

$$\text{Projection} : (\pi_{\nabla} \nabla_{\mathcal{F}})_{[X]}[Y] := [\nabla_{\mathcal{F}, X} Y]$$

Adapted differential operator :  $(\rightarrow \mathcal{D}_{\mathcal{F}}(M))$   
 $D|_U = \sum_{|\alpha| \leq k} D_{\alpha} \partial_{x^1}^{\alpha^1} \dots \partial_{x^p}^{\alpha^p} \partial_{y^1}^{\alpha^{p+1}} \dots \partial_{y^q}^{\alpha^{p+q}},$   
with  $D_{\alpha}, \alpha^1 = \dots = \alpha^p = 0$  foliated functions.

Adapted differential operator :  $(\rightarrow \mathcal{D}_{\mathcal{F}}(M))$   
 $D|_U = \sum_{|\alpha| \leq k} D_{\alpha} \partial_{x^1}^{\alpha^1} \dots \partial_{x^p}^{\alpha^p} \partial_{y^1}^{\alpha^{p+1}} \dots \partial_{y^q}^{\alpha^{p+q}},$   
with  $D_{\alpha}, \alpha^1 = \dots = \alpha^p = 0$  foliated functions.

Foliated differential operator :  $(\rightarrow \mathcal{D}(M, \mathcal{F}))$

$D|_U = \sum_{|\alpha| \leq k} D_{\alpha} \partial_{y^1}^{\alpha^1} \dots \partial_{y^q}^{\alpha^q},$   
with  $D_{\alpha}$  foliated functions.

Adapted differential operator :  $(\rightarrow \mathcal{D}_{\mathcal{F}}(M))$   
 $D|_U = \sum_{|\alpha| \leq k} D_{\alpha} \partial_{x^1}^{\alpha^1} \dots \partial_{x^p}^{\alpha^p} \partial_{y^1}^{\alpha^{p+1}} \dots \partial_{y^q}^{\alpha^{p+q}},$   
 with  $D_{\alpha}, \alpha^1 = \dots = \alpha^p = 0$  foliated functions.

Foliated differential operator :  $(\rightarrow \mathcal{D}(M, \mathcal{F}))$

$D|_U = \sum_{|\alpha| \leq k} D_{\alpha} \partial_{y^1}^{\alpha^1} \dots \partial_{y^q}^{\alpha^q},$   
 with  $D_{\alpha}$  foliated functions.

Projection  $\pi_D :$

$$\sum_{|\alpha| \leq k} D_{\alpha} \partial_{x^1}^{\alpha^1} \dots \partial_{x^p}^{\alpha^p} \partial_{y^1}^{\alpha^{p+1}} \dots \partial_{y^q}^{\alpha^{p+q}} \mapsto$$

$$\sum_{|\alpha| \leq k, \alpha^1 = \dots = \alpha^p = 0} D_{\alpha} \partial_{y^1}^{\alpha^1} \dots \partial_{y^q}^{\alpha^q}.$$

Natural  
equivariant  
quantizations

Fabian Radoux

Introduction

Cartan fiber  
bundles and  
connections

The case of the  
densities

Other differential  
operators

Conformal case

Quantizations  
and foliations

Adapted symbols : graded space  $\mathcal{S}_{\mathcal{F}}(M)$  associated to  $\mathcal{D}_{\mathcal{F}}(M)$

Adapted symbols : graded space  $\mathcal{S}_{\mathcal{F}}(M)$  associated to  $\mathcal{D}_{\mathcal{F}}(M)$

Foliated symbols : graded space  $\mathcal{S}(M, \mathcal{F})$  associated to  $\mathcal{D}(M, \mathcal{F})$



Adapted symbols : graded space  $\mathcal{S}_{\mathcal{F}}(M)$  associated to  $\mathcal{D}_{\mathcal{F}}(M)$

Foliated symbols : graded space  $\mathcal{S}(M, \mathcal{F})$  associated to  $\mathcal{D}(M, \mathcal{F})$

Projection  $\pi_{\mathcal{S}} : \pi_{\mathcal{S}}[D] = [\pi_D D]$

Adapted symbols : graded space  $\mathcal{S}_{\mathcal{F}}(M)$  associated to  $\mathcal{D}_{\mathcal{F}}(M)$

Foliated symbols : graded space  $\mathcal{S}(M, \mathcal{F})$  associated to  $\mathcal{D}(M, \mathcal{F})$

Projection  $\pi_S : \pi_S[D] = [\pi_D D]$

Conclusion :  $\pi_D(Q_{ad}(\nabla_{ad})(S_{ad})) = Q_{fol}(\pi_{\nabla}\nabla_{ad})(\pi_S S_{ad})$